

Useful Mathematical Quantities

TRIGONOMETRIC IDENTITIES

$$\cos(x \pm y) = \cos(x)\cos(y) \mp \sin(x)\sin(y) \quad (\text{C-1})$$

$$\sin(x \pm y) = \sin(x)\cos(y) \pm \cos(x)\sin(y) \quad (\text{C-2})$$

$$\cos\left(x \pm \frac{\pi}{2}\right) = \mp \sin(x) \quad (\text{C-3})$$

$$\sin\left(x \pm \frac{\pi}{2}\right) = \pm \cos(x) \quad (\text{C-4})$$

$$\cos(2x) = \cos^2(x) - \sin^2(x) \quad (\text{C-5})$$

$$\sin(2x) = 2\sin(x)\cos(x) \quad (\text{C-6})$$

$$2\cos(x) = e^{jx} + e^{-jx} \quad (\text{C-7})$$

$$2j\sin(x) = e^{jx} - e^{-jx} \quad (\text{C-8})$$

$$2\cos(x)\cos(y) = \cos(x-y) + \cos(x+y) \quad (\text{C-9})$$

$$2\sin(x)\sin(y) = \cos(x-y) - \cos(x+y) \quad (\text{C-10})$$

$$2\sin(x)\cos(y) = \sin(x-y) + \sin(x+y) \quad (\text{C-11})$$

$$2\cos^2(x) = 1 + \cos(2x) \quad (\text{C-12})$$

$$2\sin^2(x) = 1 - \cos(2x) \quad (\text{C-13})$$

$$4\cos^3(x) = 3\cos(x) + \cos(3x) \quad (\text{C-14})$$

$$4\sin^3(x) = 3\sin(x) - \sin(3x) \quad (\text{C-15})$$

$$8\cos^4(x) = 3 + 4\cos(2x) + \cos(4x) \quad (\text{C-16})$$

$$8 \sin^4(x) = 3 - 4 \cos(2x) + \cos(4x) \quad (\text{C-17})$$

$$A \cos(x) - B \sin(x) = R \cos(x + \theta) \quad (\text{C-18})$$

where

$$R = \sqrt{A^2 + B^2} \quad (\text{C-19a})$$

$$\theta = \tan^{-1}(B/A) \quad (\text{C-19b})$$

$$A = R \cos(\theta) \quad (\text{C-19c})$$

$$B = R \sin(\theta) \quad (\text{C-19d})$$

INDEFINITE INTEGRALS

Rational Algebraic Functions

$$\int (a + bx)^n dx = \frac{(a + bx)^{n+1}}{b(n+1)} \quad 0 < n \quad (\text{C-20})$$

$$\int \frac{dx}{a + bx} = \frac{1}{b} \ln |a + bx| \quad (\text{C-21})$$

$$\int \frac{dx}{(a + bx)^n} = \frac{-1}{(n-1)b(a + bx)^{n-1}} \quad 1 < n \quad (\text{C-22})$$

$$\begin{aligned} \int \frac{dx}{c + bx + ax^2} &= \frac{2}{\sqrt{4ac - b^2}} \tan^{-1} \left(\frac{2ax + b}{\sqrt{4ac - b^2}} \right) & b^2 < 4ac \\ &= \frac{1}{\sqrt{b^2 - 4ac}} \ln \left| \frac{2ax + b - \sqrt{b^2 - 4ac}}{2ax + b + \sqrt{b^2 - 4ac}} \right| & b^2 > 4ac \\ &= \frac{-2}{2ax + b} & b^2 = 4ac \end{aligned} \quad (\text{C-23})$$

$$\int \frac{x dx}{c + bx + ax^2} = \frac{1}{2a} \ln |ax^2 + bx + c| - \frac{b}{2a} \int \frac{dx}{c + bx + ax^2} \quad (\text{C-24})$$

$$\int \frac{dx}{a^2 + b^2 x^2} = \frac{1}{ab} \tan^{-1} \left(\frac{bx}{a} \right) \quad (\text{C-25})$$

$$\int \frac{x dx}{a^2 + x^2} = \frac{1}{2} \ln(a^2 + x^2) \quad (\text{C-26})$$

$$\int \frac{x^2 dx}{a^2 + x^2} = x - a \tan^{-1} \left(\frac{x}{a} \right) \quad (\text{C-27})$$

$$\int \frac{dx}{(a^2 + x^2)^2} = \frac{x}{2a^2(a^2 + x^2)} + \frac{1}{2a^3} \tan^{-1} \left(\frac{x}{a} \right) \quad (\text{C-28})$$

$$\int \frac{x dx}{(a^2 + x^2)^2} = \frac{-1}{2(a^2 + x^2)} \quad (\text{C-29})$$

$$\int \frac{x^2 dx}{(a^2 + x^2)^2} = \frac{-x}{2(a^2 + x^2)} + \frac{1}{2a} \tan^{-1}\left(\frac{x}{a}\right) \quad (\text{C-30})$$

$$\int \frac{dx}{(a^2 + x^2)^3} = \frac{x}{4a^2(a^2 + x^2)^2} + \frac{3x}{8a^4(a^2 + x^2)} + \frac{3}{8a^5} \tan^{-1}\left(\frac{x}{a}\right) \quad (\text{C-31})$$

$$\int \frac{x^2 dx}{(a^2 + x^2)^3} = \frac{-x}{4(a^2 + x^2)^2} + \frac{x}{8a^2(a^2 + x^2)} + \frac{1}{8a^3} \tan^{-1}\left(\frac{x}{a}\right) \quad (\text{C-32})$$

$$\int \frac{x^4 dx}{(a^2 + x^2)^3} = \frac{a^2 x}{4(a^2 + x^2)^2} - \frac{5x}{8(a^2 + x^2)} + \frac{3}{8a} \tan^{-1}\left(\frac{x}{a}\right) \quad (\text{C-33})$$

$$\int \frac{dx}{(a^2 + x^2)^4} = \frac{x}{6a^2(a^2 + x^2)^3} + \frac{5x}{24a^4(a^2 + x^2)^2} + \frac{5x}{16a^6(a^2 + x^2)} + \frac{5}{16a^7} \tan^{-1}\left(\frac{x}{a}\right) \quad (\text{C-34})$$

$$\int \frac{x^2 dx}{(a^2 + x^2)^4} = \frac{-x}{6(a^2 + x^2)^3} + \frac{x}{24a^2(a^2 + x^2)^2} + \frac{x}{16a^4(a^2 + x^2)} + \frac{1}{16a^5} \tan^{-1}\left(\frac{x}{a}\right) \quad (\text{C-35})$$

$$\int \frac{x^4 dx}{(a^2 + x^2)^4} = \frac{a^2 x}{6(a^2 + x^2)^3} - \frac{7x}{24(a^2 + x^2)^2} + \frac{x}{16a^2(a^2 + x^2)} + \frac{1}{16a^3} \tan^{-1}\left(\frac{x}{a}\right) \quad (\text{C-36})$$

$$\int \frac{dx}{a^4 + x^4} = \frac{1}{4a^3\sqrt{2}} \ln\left(\frac{x^2 + ax\sqrt{2} + a^2}{x^2 - ax\sqrt{2} + a^2}\right) + \frac{1}{2a^3\sqrt{2}} \tan^{-1}\left(\frac{ax\sqrt{2}}{a^2 - x^2}\right) \quad (\text{C-37})$$

$$\int \frac{x^2 dx}{a^4 + x^4} = -\frac{1}{4a\sqrt{2}} \ln\left(\frac{x^2 + ax\sqrt{2} + a^2}{x^2 - ax\sqrt{2} + a^2}\right) + \frac{1}{2a\sqrt{2}} \tan^{-1}\left(\frac{ax\sqrt{2}}{a^2 - x^2}\right) \quad (\text{C-38})$$

Trigonometric Functions

$$\int \cos(x) dx = \sin(x) \quad (\text{C-39})$$

$$\int x \cos(x) dx = \cos(x) + x \sin(x) \quad (\text{C-40})$$

$$\int x^2 \cos(x) dx = 2x \cos(x) + (x^2 - 2) \sin(x) \quad (\text{C-41})$$

$$\int \sin(x) dx = -\cos(x) \quad (\text{C-42})$$

$$\int x \sin(x) dx = \sin(x) - x \cos(x) \quad (\text{C-43})$$

$$\int x^2 \sin(x) dx = 2x \sin(x) - (x^2 - 2) \cos(x) \quad (\text{C-44})$$

Exponential Functions

$$\int e^{ax} dx = \frac{e^{ax}}{a} \quad a \text{ real or complex} \quad (\text{C-45})$$

$$\int x e^{ax} dx = e^{ax} \left[\frac{x}{a} - \frac{1}{a^2} \right] \quad a \text{ real or complex} \quad (\text{C-46})$$

$$\int x^2 e^{ax} dx = e^{ax} \left[\frac{x^2}{a} - \frac{2x}{a^2} + \frac{2}{a^3} \right] \quad a \text{ real or complex} \quad (\text{C-47})$$

$$\int x^3 e^{ax} dx = e^{ax} \left[\frac{x^3}{a} - \frac{3x^2}{a^2} + \frac{6x}{a^3} - \frac{6}{a^4} \right] \quad a \text{ real or complex} \quad (\text{C-48})$$

$$\int e^{ax} \sin(x) dx = \frac{e^{ax}}{a^2 + 1} [a \sin(x) - \cos(x)] \quad (\text{C-49})$$

$$\int e^{ax} \cos(x) dx = \frac{e^{ax}}{a^2 + 1} [a \cos(x) + \sin(x)] \quad (\text{C-50})$$

DEFINITE INTEGRALS

$$\int_{-\infty}^{\infty} e^{-a^2 x^2 + bx} dx = \frac{\sqrt{\pi}}{a} e^{b^2/(4a^2)} \quad a > 0 \quad (\text{C-51})$$

$$\int_0^{\infty} x^2 e^{-x^2} dx = \sqrt{\pi}/4 \quad (\text{C-52})$$

$$\int_0^{\infty} \text{Sa}(x) dx = \int_0^{\infty} \frac{\sin(x)}{x} dx = \frac{\pi}{2} \quad (\text{C-53})$$

$$\int_0^{\infty} \text{Sa}^2(x) dx = \pi/2 \quad (\text{C-54})$$

$$\sum_{n=1}^N n = \frac{N(N+1)}{2} \quad (\text{C-55})$$

$$\sum_{n=1}^N n^2 = \frac{N(N+1)(2N+1)}{6} \quad (\text{C-56})$$

$$\sum_{n=1}^N n^3 = \frac{N^2(N+1)^2}{4} \quad (\text{C-57})$$

$$\sum_{n=0}^N x^n = \frac{x^{N+1} - 1}{x - 1} \quad (\text{C-58})$$

$$\sum_{n=0}^N \frac{N!}{n!(N-n)!} x^n y^{N-n} = (x+y)^N \quad (\text{C-59})$$

$$\sum_{n=0}^N e^{j(\theta+n\phi)} = \frac{\sin[(N+1)\phi/2]}{\sin(\phi/2)} e^{j[\theta+(N\phi/2)]} \quad (\text{C-60})$$

$$\sum_{n=0}^N \binom{N}{n} = \sum_{n=0}^N \frac{N!}{n!(N-n)!} = 2^N \quad (\text{C-61})$$

$$\sum_{n=N_1}^{N_2} w^n = \frac{w^{N_1} + w^{N_2+1}}{1-w} \quad \begin{cases} N_2 > N_1 \text{ and } w \\ \text{real or complex} \end{cases} \quad (\text{C-62})$$

INFINITE SERIES

$$e^x = 1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} + \cdots = \sum_{n=0}^{\infty} \frac{x^n}{n!} \quad (\text{C-63})$$